


Nafith Abu Tarboush
DDS, MSc, PhD
natarboush@ju.edu.jo
www.facebook.com/natarboush

Oxidative Phosphorylation




The Mitochondria

- OMM: permeable to small molecules (**MW < 5,000**) & ions, porins (transmembrane channels)
- IMM: impermeable even to H^+ ; specific transporters
- IMM bears the components of the respiratory chain and the ATP synthase
- Matrix: contains pyruvate dehydrogenase complex & TCA cycle enzymes, fatty acid β -oxidation pathway, and the pathways of amino acid oxidation
- In other words: matrix contains all pathways of fuel oxidation except glycolysis (cytosol)




The oxidative phosphorylation, Where are we?

➤ Stages: Digestion; Acetyl-CoA, TCA, OxPhos




Oxidative phosphorylation (OxPhos)

- Generation of ATP aided by the reduction of O_2
- Peter Mitchell (1961): the chemiosmotic theory
- Oxidative phosphorylation have 3 major aspects:
 - ✓ (1) It involves flow of electrons through a chain of membrane-bound carriers (prosthetic groups)
 - ✓ (2) The free energy available (exergonic) is coupled to transport protons across a proton-impermeable membrane
 - ✓ (3) The transmembrane flow of protons down their concentration gradient provides the free energy for synthesis of ATP (ATP synthase)






Oxidative phosphorylation (OxPhos)



Types of electron transfer (ET) through the electron transport chain (ETC)

- 3 types of ET occur in OxPhos:
 - ✓ Direct ET, as in the reduction of Fe^{+3} to Fe^{+2}
 - ✓ Transfer as a hydrogen atom $\{(\text{H}^+) + (\text{e}^-)\}$
 - ✓ Transfer as a hydride ion $(:\text{H}^-)$








Electrons are funneled to a universal electron acceptors

COENZYME	AS OXIDIZING AGENT	AS REDUCING AGENT
Nicotinamide adenine dinucleotide	NAD^+	NADH/H^+
Nicotinamide adenine dinucleotide phosphate	NADP^+	NADPH/H^+
Flavin adenine dinucleotide	FAD	FADH_2
Flavin mononucleotide	FMN	FMNH_2

Other electron-carrying molecules

“Ubiquinone”



- Also called coenzyme Q, or Q
- Lipid-soluble benzoquinone with a long isoprenoid side chain
- Small & hydrophobic (freely diffusible)
- Carries electrons through the IMM
- Can accept either 1 e- or 2 e-
- Act at the junction between a 2-electron donor and a 1-electron acceptor
- Sometimes prescribed for recovering MI patients



Other electron-carrying molecules


“Cytochromes”

- Proteins with characteristic strong absorption of visible light (Fe-containing heme prosthetic groups)
- Classification based on light absorption
- Mode of binding (a, b, c)
- Mitochondria contain three classes of cytochromes (a, b, & c)




Requirements of OxPhos

- Redox reaction: electron donor (NADH or FADH₂) & electron acceptor (O₂)
- An intact IMM
- ETC of proteins
- ATP synthase



ET to O₂, how does the process occurs? “The chemi-osmotic theory”



- Membrane is impermeable to protons
- ≈ - 16 kcal with each step

Oxi-Red Components of ETC







➤ "NADH Dehydrogenase" – Complex I

➤ NADH-Q oxidoreductase


➤ "Succinate Dehydrogenase" – Complex II

➤ ≈ 0 kcal, H^+ ?

➤ "Cytochrome bc₁" – Complex III
Q-cytochrome c Oxidoreductase




"Cytochrome c oxidase" – Complex IV



Pumping of Protons

Intermembrane space (P side)






➤ For every 2 electrons passing:

➤ 4H^+ (complex I); 0H^+ (complex II); 4H^+ (complex III), 2H^+ (complex IV)


ATP Synthase

- **F₁:**
 - “γ” subunit: rotates
 - “β” subunit: binds
 - “α” subunit: structural
 - 3 conformations: tight (T), loose (L), open (O)
- **F₀:**
 - “a” subunit: point of entry & exit
 - “c” subunit rotates
 - 4H⁺/ATP
 - Can run backwards



Energy yield from the ETC

- NADH, -53 kcal, ATP?
- FADH₂, -41 kcal, ATP?
- ΔG° is so negative, never reversible
- ATP machine efficiency, (anions, Ca²⁺, heat, phosphate, substrates)
- Electron transport chain is our major source of heat




Regulation – the need for ATP

- What OxPhos needs? (NADH, O₂, ADP, and Pi)
- In skeletal muscles, 20% drop in ATP concentration
- In the heart, Ca⁺² activates TCA enzymes for extra push (NADH, ATP), no drop
- ET is tightly coupled to phosphorylation (simultaneously)
- ADP is the most important factor in determining the rate
- The regulation of the rate of oxidative phosphorylation by the ADP level is called respiratory control




Regulation – inhibition (coupling)

- Can occur at any stage
- Specific inhibitors:
 - ✓ Cyanoglycosides such as amygdalin are present in edible plant pits
 - ✓ Oligomycin prevents the influx of H⁺ through ATP synthase (tight coupling)



Anit-cancerous drug

Specific inhibitor	Target
Rotenone (insecticide) & Amytal (sedative)	NADH-Q oxidoreductase
Antimycin A (antibiotic)	Q-cytochrome c oxidoreductase
Cyanide (CN ⁻), Azide (N ₃ ⁻), & (CO)	Cytochrome c oxidase
Oligomycin (antibiotic)	ATP synthase



الصفحة الرئيسية > محليات



أشهر جرائم القتل العائلية في المملكة

جراسا نيوز -

جراسا - تعرض فيما يلي قائمة بأشهر جرائم القتل العائلية التي حدثت في الأردن خلال السنوات الماضية ، والتي كان لكل منها وقع الصدمة حين وقوعها لما تمثله من فعل غريب على المجتمع وأعرافه ، فضلا عن مخالفتها للشرائع السماوية والقوانين النافذة والطبيعة الإنسانية بعامه.

قضية السيائيد

أول جريمة من نوعها يرتكبها أب ضد ولديه ، إذ قام الأب بوضع مادة السيائيد في كأس الحليب وطلب من طفليه ان يشربا منه ، حيث فارقا الحياة بعد 10 دقائق من مغادرة الام المنزل لتعود وتجدهما جثتين هامدتين.



وقد ادبى الاب بعقوبة الاعدام شتفا الا ان والده اسقط الحق الشخصي كونه وليا عن الطفلين وحكم عليه بالاشغال المؤبدة.



Regulation – Uncoupling

Unregulated – chemical uncouplers



- What is uncoupling?
- How does it occur? Dissipation of PMF
- What is the result?
- Is it physiological or not?
- 2,4-dinitrophenol (DNP) & other acidic aromatic compounds
- What changes happen? \uparrow O_2 consumption, \uparrow NADH oxidation
- Soviet soldiers were given DNP, FDA banned DNP (1938)



Regulation – Uncoupling

Regulated - Uncoupling proteins (UCPs)

- Short-circuiting ATP synthase
- UCP₁ (thermogenin):
 - ✓ Brown adipose tissue, non-shivering thermogenesis
 - ✓ Infants: neck, breast, around kidneys
 - ✓ Fatty acids directly activates UCP₁
- UCP₂ (most cells); UCP₃ (skeletal muscle); {UCP₄, UCP₅} (brain)
- Obesity tendency in some populations




OxPhos Diseases (Genetic)

- **A. Mitochondrial DNA and OXPHOS Diseases**
 - ✓ **Small (16,569) base pair, double-stranded, circular DNA**
 - ✓ **Encodes 13 subunits: 7 (I) , 1 (III), 3 (IV), 2 (Fo)**
 - ✓ **Also encodes necessary components for translation of its own mRNA: a large and small rRNA and tRNAs**
 - ✓ **Maternal inheritance, heteroplasmy**
 - ✓ **Accumulation of somatic mutations with age**
 - ✓ **Highest ATP demands: CNS, heart, skeletal muscle, and kidney, liver**

OxPhos Diseases (Genetic)

- **B. Nuclear Genetic Disorders of Oxidative Phosphorylation**
 - ✓ **1,000 proteins (50% of the mitochondria is protein)**
 - ✓ **Usually autosomal recessive**
 - ✓ **Expressed in all tissues**
 - ✓ **Phenotypic expression with high ATP demand**



OxPhos Diseases (Genetic)



Mitochondrial shuttling systems

“Cytosolic NADH”


- Glycerol 3-phosphate shuttle (Sk. muscles & brain)
- Glycolytic pathway as an example
- How NADH passes?
- ATP yield?



Mitochondrial shuttling systems

“Cytosolic NADH”

- Malate-Aspartate shuttle
- Heart & liver
- 2 membrane carriers & 4 enzymes
- Readily reversible (vs. Glycerol 3-phosphate shuttle)
- NADH can be transferred only if the NADH/NAD⁺ ratio is higher in the cytosol than in the mitochondrial matrix
- Exchange of key intermediates between mitochondria & cytosol



Mitochondrial shuttling systems

“ATP/ADP”

➤ ATP-ADP Translocase (also called adenine nucleotide translocase or ANT)

➤ The flows of ATP and ADP are coupled (ADP enters only if ATP exits, and vice versa)

➤ Highly abundant (14% of IMM proteins)

➤ Contains a single nucleotide-binding site (alternates)

➤ Similar affinity to ATP and ADP

➤ Endergonic (25% of ETC)

➤ Inhibition leads to subsequent inhibition of cellular respiration

