

Valves of the heart

Two atrioventricular valves

Right atrioventricular
or tricuspid valve

Left atrioventricular
or bicuspid valve
Mitral valve

Two semilunar valves

Aortic valve

Pulmonary valve

The main function of any valve is to maintain the Unidirectional flow of blood


RIGHT ATRIOVENTRICULAR VALVE

The TRICUSPID valve


so-named because it usually consists of *three cusps or leaflets*

The bases of the valve cusps are attached to the fibrous ring around the orifice ***annulus fibrosus***

- Its orifice is best seen from the atrial aspect and measures on average **11.4 cm**
- The fibrous ring keeps the caliber of the orifice constant (large enough to admit the tips of three fingers)
- It is almost **VERTICAL!!!!!!!!!!**, but at 45° to the sagittal plane
- The atrial surface of the AV valve is rather smooth.
- The ventricular surface is irregular because of the insertion of the chordae tendineae



guards the right atrioventricular orifice



It is almost VERTICAL

**Heart in diastole:
viewed from base with atria removed**

The atrioventricular valvular complex

Consists of :

- 1- *The orifice and its associated anulus*
- 2- *The cusps*, the supporting *chordae tendineae* of various types and *the papillary muscles*




Fig. 56.10 Principal elements of the fibrous skeleton of the heart. For clarity, the view is from the right posteriosuperior aspect. Perspective causes the pulmonary anulus to appear smaller than the aortic anulus, whereas in fact the reverse is the case. Consult text for an extended discussion. (Copyright from The Royal College of Surgeons of England. Reproduced with permission.)

Anterior

➤ The Tricuspid Valve consists of three cusps



PULMONARY VALVE

Right semilunar cusp

Anterior semilunar cusp

Left semilunar cusp

AORTIC VALVE

Left semilunar cusp

Right semilunar cusp

Posterior semilunar cusp

L

MITRAL VALVE

Anterior cusp

Posterior cusp

Anterior cusp lies anteriorly

Septal cusp lies against the ventricular septum

inferior (posterior) cusp lies inferiorly

TRICUSPID VALVE


Anterior cusp

Posterior cusp

Septal cusp


A.Sk

Posterior




- The bases of the cusps are attached to the fibrous ring of the skeleton of the heart whereas their free edges and ventricular surfaces are attached to the chordae tendineae.

- The chordae tendineae connect the cusps to the papillary muscles



Tricuspid (right atrioventricular) valve

The Tricuspid Valve



The cusps



The chordae tendineae.

papillary muscles

Cusp of tricuspid valve

Chordae tendineae

Papillary muscle



a

b

Histology

Valves **Cusps** are composed of connective tissue with over-lying endocardium.

Each valve is composed of three layers


The spongiosa is loose connective tissue located on the **atrial or blood vessel side (aortic and pulmonary)** of each valve.

The fibrosa forms the core of the valve and contains **fibrous extensions from the dense irregular connective tissue of the skeletal rings of the heart**

The ventricularis is immediately adjacent to the ventricular surface of each valve and is covered with **endothelium**.

It contains **dense connective tissue with many layers of elastic fibers**.

In the AV valves, the ventricularis continues into **the chordae tendineae**



Valve cusps are normally avascular

Small blood vessels and smooth muscle can be found only in the base of the cusp. The surfaces of the valve are exposed to blood, and the cusps are thin enough to allow nutrients and oxygen to diffuse from the blood



Normal conditions

(valvulitis).

- Rheumatic fever causes inflammation of the heart valves (valvulitis)
- Inflammation induces **angiogenesis in the valve and vascularization** in the normally avascular layers of the valve.
 - These changes most commonly affect the **mitral valve (65% to 70%)** ?????!! and aortic valve (20% to 25%).
- This inflammation can lead to ***progressive replacement of elastic tissue by irregular masses of collagen fibers***, causing the valve **to thicken**.
- The valves become rigid and inflexible, which **affects their ability to open and close**

The extreme edges of the cusps are thin and delicate with a **sawtooth appearance** from the insertion of **chordae**

Away from the edge, the atrial surface of the cusps is finely **nodular**, the nodule particularly in small children. These nodules are called **The noduli Albin**



The noduli Albini


minute fibrous nodules on the margins of the mitral and tricuspid valves of the heart

Fibrous (Albini's) nodules

When the ventricle contracts, **the papillary muscles contract** and *prevent the cusps* from being forced into the atrium and turning inside out as the intraventricular pressure rises

To assist in this process, the chordae tendineae of one papillary muscle are connected to the **adjacent parts of two cusps**

On closure of an AV valve, the narrow border between the **row of Albini nodules** and the free edge of each cusp presses against that of the next, resulting in a secure, **watertight closure**



Tricuspid (right atrioventricular) valve

Posterior

Tricuspid valve

Bicuspid (mitral) valve

A.S.H


Right side of heart

Left side of heart

Aortic valve

Pulmonary valve

Anterior





Chordae tendineae (tendinous cords)

are fibrous collagenous structures supporting the cusps of the atrioventricular valves

Tendinous cords attach to the free edges and ventricular surfaces of the anterior, posterior, and septal cusps, much like the cords attaching to a parachute

The tendinous cords arise from the apices of papillary muscles



Papillary muscles begin to contract before contraction of the right ventricle, tightening the tendinous cords and drawing the cusps together.

Because the cords are **attached to adjacent sides of two cusps**,



they prevent separation of the cusps and

prevented from prolapsing (being driven into the right atrium) as ventricular pressure rises.


Thus, **regurgitation of blood** (backward flow of blood) from the right ventricle back into the right atrium is blocked during ventricular systole by the valve cusp

Do papillary muscles close valve or do they open it?


Papillary muscles insure Competence of the valve



The mitral valve



- guards the left atrioventricular orifice
- It consists of two cusps.






Fig. 56.13 The aortic orifice opened from the front to show the cusps of the aortic valves, their nodules, lunules, commissures and the triple-scalloped line of anular attachment. Also shown is the continuity of the subaortic curtain with the mitral anterior cusp (i.e. 'aortic baffle') and the coronary ostia, and the spatial relationship of the aortic orifice to the pulmonary orifice and to the left ventricle. (From Sobotta 2006.)


The anterior cusp of the mitral valve

THE SUBAORTIC CURTAIN


Subjected to blood flow from two sides ???

- The left atrioventricular orifice admits atrial blood during diastole, flow being towards the cardiac apex.
 - After closure of the mitral cusps, and throughout the ejection phase of systole, blood is expelled from the apex through the aortic orifice




The bloodstream undergoes two right angle turns, which together result in a 180° change in direction. This reversal of flow takes place around **the anterior cusp of the mitral valve**



Semilunar valves



No chordae or papillary muscles are associated with these valves



The pulmonary valve guards the pulmonary orifice and

The aortic valve guards the aortic orifice.

Each consists of three pocketlike cusps of approximately equal size


There is no distinct, circular ring of fibrous tissue at the base of the arteries from which these and the valve cusps arise

The arterial wall **expands into three dilated pouches**, the sinuses of **Valsalva**

The cusps of the arterial semilunar valve are largely smooth and thin.

At the center of the free margin of each cusp is a small fibrous nodule called

The nodulus Arantii



On each side of the nodules of Arantius, along the entire free edge of the cusp, there is a thin, halfmoon-shaped area called the **lunula** that has fine striations parallel to the edge

The aortic valve


consists of three semilunar cusps

Posterior (non-coronary) cusp

Right

Left

Just superior to right and left cusps in the Sinus of **Valsalva** are the openings of the right and left coronary arteries, respectively



(A) Anterior view of aortic valve

(B) Valve open

(C) Valve closed

Pulmonary valve

At the apex of the infundibulum, the outflow tract of the right ventricle, the opening into the pulmonary trunk is closed by the **pulmonary valve**

consists of three **semilunar cusps** with free edges projecting upward into the lumen of the pulmonary trunk

3 semilunar cusps

Anterior

Right

Left

Opening of the pulmonary valve

During diastole, the pulmonary valve is closed and all three cusps of the valve are tightly apposed.

The pulmonary valve opens passively during ventricular systole and then closes rapidly at the end of systole

