

# Pneumonia

DR. NASER MAHMOUD


# Pneumonia

- ▶ Definitions
- ▶ Classifications
- ▶ Epidemiology
- ▶ Pathophysiology/Pathogenesis
- ▶ Microbiology
- ▶ History
- ▶ Physical Examination
- ▶ Investigations: Imaging and labs
- ▶ Complications
- ▶ Differential diagnosis
- ▶ Management
- ▶ Prevention

# Definitions

- ▶ The term **"Pneumonia"** is usually used to describe an **infection** of the lung **parenchyma**. Although it is still used to describe some non infectious diseases (e.g. eosinophilic pneumonia).
- ▶ Don't confuse it with other infections of the lower respiratory tract.
- ▶ Because it is affecting the lung parenchyma almost all pneumonias will have some **radiologic findings** (with rare exceptions)


# Lower respiratory and pleural disease



**Empyema:** purulent exudate in the pleural cavity

**Abscess:** circumscribed collection of pus within the lung parenchyma

**Pneumonia** -- infection of alveoli (viral or bacterial)  
vs. **Pneumonitis** -- immune-mediated inflammation of alveoli, XRT pneumonitis, aspiration pneumonitis...



**Bronchitis** -- inflammation of bronchi, may be **immune-mediated**, e.g. asthma, COPD, or **infectious** (usually viral but can be bacterial)



**Bronchiolitis:** inflammation of bronchioles (often viral but can be bacterial or autoimmune)

# Classifications

ATS/IDSA:

- ▶ Community Acquired Pneumonia
- ▶ Hospital Acquired Pneumonia
- ▶ Ventilator Associated Pneumonia
- ▶ Others (Pneumonia in immunocompromised patients, TB etc..)

What are these:


- ▶ HCAP???
- ▶ Atypical??
- ▶ Pneumonia based on organisms??
  
- ▶ Why it is important to know this classification??

# Epidemiology

- ▶ Accounts for around 10% of all admissions to hospitals in the united states.
- ▶ A significant cause of morbidity and mortality in adults.

## Top 10 Causes of Death in the USA


(according to Centers for Disease Control and Prevention)



passare.com

Twitter: @PassareInc  
Facebook: facebook.com/passareinc  
Tumblr: passareinc.tumblr.com


## Top 10 causes of deaths in low-income countries in 2016



Source: Global Health Estimates 2016: Deaths by Cause, Age, Sex, by Country and by Region, 2000-2016. Geneva, World Health Organization; 2018.  
World Bank list of economies (June 2017). Washington, DC: The World Bank Group; 2017 (<https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups>).

## Top 10 causes of deaths in upper-middle-income countries in 2016


Crude death rate (per 100 000 population)



Source: Global Health Estimates 2016: Deaths by Cause, Age, Sex, by Country and by Region, 2000-2016. Geneva, World Health Organization; 2018.  
World Bank list of economies (June 2017). Washington, DC: The World Bank Group; 2017 (<https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups>).

## Top 10 causes of deaths in lower-middle-income countries in 2016

Crude death rate (per 100 000 population)



Source: Global Health Estimates 2016: Deaths by Cause, Age, Sex, by Country and by Region, 2000-2016. Geneva, World Health Organization; 2018.  
World Bank list of economies (June 2017). Washington, DC: The World Bank Group; 2017 (<https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups>).

# Pathophysiology/Pathogenesis

**Is the lower respiratory tract a sterile area???**

**Interactions between host and pathogens.**

- ▶ **Inhalation**
- ▶ **Aspiration**
- ▶ **Hematogenous spread**
- ▶ **Direct extension from adjacent infected foci**
- ▶ Tuberculosis can spread contiguously from the lymph nodes to the pericardium or the lung, but this is only rarely a route of pneumonia formation

# Microbiology

Mild to severe

<ol style="list-style-type: none"><li>1. <i>Mycoplasma pneumoniae</i></li><li>2. <i>Streptococcus pneumoniae</i></li><li>3. <i>Chlamydomphila pneumoniae</i></li><li>4. <i>Haemophilus influenzae</i></li><li>5. Respiratory viruses</li></ol>	<ol style="list-style-type: none"><li>1. <i>Streptococcus pneumoniae</i></li><li>2. <i>Mycoplasma pneumoniae</i></li><li>3. <i>Chlamydomphila pneumoniae</i></li><li>4. <i>Haemophilus influenzae</i></li><li>5. <i>Staphylococcus aureus</i></li><li>6. Mixed infections</li><li>7. Enteric gram-negative bacilli</li><li>8. Aspiration (anaerobes)</li><li>9. Respiratory viruses</li><li>10. <i>Legionella</i> species</li></ol>	<ol style="list-style-type: none"><li>1. <i>Streptococcus pneumoniae</i></li><li>2. Enteric gram-negative bacilli</li><li>3. <i>Staphylococcus aureus</i></li><li>4. <i>Legionella</i> species</li><li>5. <i>Mycoplasma pneumoniae</i></li><li>6. Respiratory viruses</li><li>7. <i>Pseudomonas aeruginosa</i> (relative frequency determined by the presence or absence of specific risk factors)</li></ol>
--	---	--

This is in the general population. In special patients groups like immunocpmpromised patients other pathogens can cause pneumonia (e.g. fungal infections, CMV, Atypical mycobacterium...etc)

# History

- ▶ Pneumonia is characterized by the presence of fever, altered general well-being, and respiratory symptoms, such as cough (90%), sputum production (66%), dyspnea (66%), pleuritic pain (50%), and hemoptysis (15%).
- ▶ In older and immunocompromised patients, the signs and symptoms of pulmonary infection may be muted and overshadowed by nonspecific complaints, pneumonia may present with general weakness, decreased appetite, altered mental status, incontinence, or decompensation due to underlying disease.
- ▶ Pneumococcal pneumonia: “classic” history, such as that of the patient with pneumococcal infection who presents with sudden onset of rigor followed by pleuritic chest pain, dyspnea, and cough with rusty sputum.
- ▶ *Legionella* pneumonia may complain predominantly of diarrhea, fever, headache, confusion, and myalgia.
- ▶ *M. pneumoniae* infection, extrapulmonary manifestations such as myringitis, encephalitis, uveitis, iritis, and myocarditis may be present. However, only rarely does the clinical history clearly suggest a specific etiologic diagnosis.
- ▶ **Information obtained from the clinical history and physical examination is not sufficient to confirm the diagnosis of pneumonia. A definitive diagnosis requires the finding of a new opacity on the chest radiograph.**



- ▶ **Typical Vs. Atypical syndromes.**

- ▶ “Typical” characteristically caused by bacteria such as *S. pneumoniae*, *H. influenzae*, and *K. pneumoniae*. The initial presentation is frequently acute, with an intense chill. Productive cough is present, and the sputum is purulent or bloody. Physical examination reveals typical findings of pulmonary consolidation . Blood tests show leukocytosis with neutrophilia and the presence of band forms in most cases. Chest radiography shows lobar consolidation with air bronchograms.

- 
- ▶ “Atypical pneumonia.” : gradual onset of fever, nonproductive cough, and a relatively normal white blood cell count in a patient without a readily demonstrable bacterial pathogen, systemic complaints are more prominent than the respiratory ones. The atypical syndrome is characteristic of infections by pathogens such as *M. pneumoniae*, *Chlamydia* species, *C. burnetii*, and viruses.
  - ▶ Neither the clinical symptoms nor the radiographic manifestations are sufficiently sensitive or specific to guide pathogen-directed antibiotic treatment against “typical” versus “atypical” microorganisms.

**TABLE 10. Possible Microbial Causes of Community-Acquired Pneumonia**

Characteristics	Commonly Encountered Pathogens
<b>Clinical Presentation</b>	
Aspiration	Gram-negative enteric pathogens, oral anaerobes
Cough >2 weeks with whoop or posttussive vomiting	<i>Bordetella pertussis</i>
Lung cavity infiltrates	Community-associated methicillin-resistant <i>Staphylococcus aureus</i> , oral anaerobes, endemic fungal pathogens, <i>Mycobacterium tuberculosis</i> , atypical mycobacteria
<b>Epidemiology or Risk Factor</b>	
Alcoholism	<i>Streptococcus pneumoniae</i> , oral anaerobes, <i>Klebsiella pneumoniae</i> , <i>Acinetobacter</i> species, <i>M. tuberculosis</i>
COPD and/or smoking	<i>Haemophilus influenzae</i> , <i>Pseudomonas aeruginosa</i> , <i>Legionella</i> species, <i>S. pneumoniae</i> , <i>Moraxella catarrhalis</i> , <i>Chlamydophila pneumoniae</i>
Exposure to bat or bird droppings	<i>Histoplasma capsulatum</i>
Exposure to birds	<i>Chlamydophila psittaci</i> (if poultry: avian influenza)
Exposure to rabbits	<i>Francisella tularensis</i>
Exposure to farm animals or parturient cats	<i>Coxiella burnetii</i>
Exposure to rodent excreta	Hantavirus
HIV infection (early)	<i>S. pneumoniae</i> , <i>H. influenzae</i> , <i>M. tuberculosis</i>
HIV infection (late)	<i>S. pneumoniae</i> , <i>H. influenzae</i> , <i>M. tuberculosis</i> , <i>Pneumocystis jirovecii</i> , <i>Cryptococcus</i> species, <i>Histoplasma</i> species, <i>Aspergillus</i> species, atypical mycobacteria (especially <i>Mycobacterium kansasii</i> ), <i>P. aeruginosa</i>
Hotel or cruise ship stay in previous 2 weeks	<i>Legionella</i> species
Travel or residence in southwestern United States	<i>Coccidioides</i> species, hantavirus
Travel or residence in Southeast and East Asia	<i>Burkholderia pseudomallei</i> , avian influenza, severe acute respiratory syndrome–coronavirus (SARS-CoV)
Travel or residence in (or exposure to an ill traveler from) the Middle East	Middle East respiratory syndrome–coronavirus (MERS-CoV)
Influenza activity in community	Influenza, <i>S. pneumoniae</i> , <i>S. aureus</i> , <i>H. influenzae</i>
Injection drug use	<i>S. aureus</i> , anaerobes, <i>M. tuberculosis</i> , <i>S. pneumoniae</i>
Endobronchial obstruction	Anaerobes, <i>S. pneumoniae</i> , <i>H. influenzae</i> , <i>S. aureus</i>
Bronchiectasis or cystic fibrosis	<i>Burkholderia cepacia</i> , <i>P. aeruginosa</i> , <i>S. aureus</i>
Bioterrorism	<i>Bacillus anthracis</i> , <i>Yersinia pestis</i> , <i>Francisella tularensis</i>

Adapted with permission from Mandell LA, Wunderink RG, Anzueto A, et al; Infectious Diseases Society of America; American Thoracic Society. Infectious Diseases Society of America/American Thoracic Society consensus guidelines on the management of community-acquired pneumonia in adults. Clin Infect Dis. 2007 Mar 1;44(suppl 2):S27-72. [PMID: 17278083] Copyright 2007, Oxford University Press.

# Physical examination

- ▶ General
- ▶ Vital signs
- ▶ Hands
- ▶ H&N
- ▶ Chest
- ▶ Heart
- ▶ Abdomen
- ▶ LEs

- 
- ▶ Expansion: reduced on the affected side.
  - ▶ Vocal fremitus: increased on the affected side (in other chest disease this sign is of very little use!).
  - ▶ Percussion: dull, but not stony dull.
  - ▶ Breath sounds: bronchial.
  - ▶ Additional sounds: medium, late or pan-inspiratory crackles as the pneumonia resolves.
  - ▶ Vocal resonance: increased.
  - ▶ Pleural rub: may be present

<b>Disorder</b>	<b>Mediastinal displacement</b>	<b>Chest wall movement</b>	<b>Percussion note</b>	<b>Breath sounds</b>	<b>Added sounds</b>
Consolidation	None	Reduced over affected area	Dull	Bronchial	Crackles
Collapse	Ipsilateral shift	Decreased over affected area	Dull	Absent or reduced	Absent
Pleural effusion	Heart displaced to opposite side (trachea displaced only if massive)	Reduced over affected area	Stony dull	Absent over fluid; may be bronchial at upper border	Absent; pleural rub may be found above effusion
Pneumothorax	Tracheal deviation to opposite side if under tension	Decreased over affected area	Resonant	Absent or greatly reduced	Absent
Bronchial asthma	None	Decreased symmetrically	Normal or decreased	Normal or reduced	Wheeze
Interstitial pulmonary fibrosis	None	Decreased symmetrically (minimal)	Normal unaffected by cough or posture	Normal	Fine, late or pan-inspiratory crackles over affected lobes

ELSEVIER

## Coarse crackles

J. Lucian Davis, MD and John F. Murray, MD

©2016 Elsevier. All Rights Reserved.

ELSEVIER

## Bronchial breath sounds

J. Lucian Davis, MD and John F. Murray, MD

©2016 Elsevier. All Rights Reserved.

ELSEVIER

## **Pleural friction rub**

J. Lucian Davis, MD and John F. Murray, MD

©2016 Elsevier. All Rights Reserved.

- ▶ **Clues to the etiologic diagnosis may lie outside the respiratory tract.**
- ▶ Bradycardia in relation to the amount of fever (pulse should increase by 10 beats/min/°C of temperature elevation) has been associated with pneumonia due to Legionella, C. psittaci, Mycoplasma, or F. tularensis.
- ▶ M. pneumoniae infection may present with extrapulmonary manifestations including arthralgia, cervical lymphadenopathy, bullous myringitis, diarrhea, myalgia, myocarditis, hepatitis, nausea, pericarditis, and vomiting. Skin lesions of erythema multiforme or erythema nodosum suggest Mycoplasma infection (as well as tuberculosis and endemic fungal infection), whereas lesions of ecthyma gangrenosum are most often seen with P. aeruginosa infection.
- ▶ Finally, the examiner must look for the presence of complications such as pleural effusion, pericarditis, endocarditis, arthritis, and central nervous system involvement, which may necessitate further diagnostic procedures and, potentially, a change in therapy.

# Laboratory Evaluation/basic labs

- ▶ CBC
- ▶ Blood Gas
- ▶ CRP
- ▶ ESR
- ▶ PCT
- ▶ KFT
- ▶ LFT

# Laboratory Evaluation/Microbiologic eval.

## PATIENTS WHO DO NOT REQUIRE HOSPITALIZATION

None\*

## PATIENTS WHO REQUIRE HOSPITALIZATION

1. Two sets of blood cultures (obtained prior to antibiotics)
2. Gram stain and culture of a valid sputum sample
3. Urinary antigen test for detection of *Legionella pneumophila* (in endemic areas or during outbreaks)
4. Stain for acid-fast bacilli and culture of sputum (if tuberculosis is suggested by clinical history or radiologic findings)
5. Fungal stain and culture of sputum, and fungal serologies (if infection by an endemic mycosis is suggested by the clinical history or radiologic findings)
6. Sputum examination for *Pneumocystis jirovecii* (if suggested by clinical history or radiologic findings)
7. Nucleic acid amplification tests for *Mycoplasma pneumoniae*, *Chlamydophila pneumoniae*, *Chlamydophila psittaci*, *Coxiella burnetii*, *Legionella* species, and respiratory viruses (in endemic areas or during outbreaks)
8. Culture and microscopic evaluation of pleural fluid (if significant fluid is present)

## ADDITIONAL TESTS FOR PATIENTS WHO REQUIRE TREATMENT IN AN ICU

1. Gram stain and culture of endotracheal aspirate or bronchoscopically obtained specimens using a protected specimen brush or BAL
2. Other procedures as for other hospitalized patients






BAL, bronchoalveolar lavage; ICU, intensive care unit.

# Clinical Indications for More Extensive Testing in Community-Acquired Pneumonia

- 1.Intensive care unit admission
- 2.Failure of outpatient antibiotic therapy
- 3.Radiographic cavities
- 4.Leukopenia
- 5.Active alcohol abuse
- 6.Chronic severe liver disease
- 7.Severe obstructive/structural lung disease
- 8.Asplenia
- 9.Recent travel (within past 2 weeks)
- 10.Positive *Legionella* UAT result
- 11.Positive pneumococcal UAT result
- 12.Pleural effusion

# Laboratory Evaluation/Sputum Examination

- ▶ Easy to do.
- ▶ A specimen with few or no squamous cells and many polymorphonuclear white blood cells (>25 cells/low-power field in a sample from a patient who is not
- ▶ The latest IDSA/ATS guidelines recommend obtaining a sputum sample for Gram stain and culture in hospitalized patients.
- ▶ In ventilated patients, the equivalent of sputum is the endotracheal aspirate.
- ▶ Some bacterial agents of pneumonia cannot be cultivated on conventional laboratory media. For example, *Legionella* requires buffered charcoal yeast extract agar for isolation, whereas recovery of *Chlamydomphila* species and *C. burnetii* requires culture in mammalian cell lines.



# Laboratory Evaluation/Blood and Pleural Fluid Cultures

- ▶ The overall yield of blood cultures is less than 20% in patients hospitalized for CAP.
- ▶ The detected rate of bacteremia is lower in patients with mild CAP and higher in patients with severe CAP, especially those warranting ICU care. Prior antibiotic treatment decreases the yield of blood cultures.
- ▶ In up to 40% of CAP cases, a pleural effusion may be present. Although the specificity of pleural exudate cultures is very high, the sensitivity is low because of the low incidence of invasion of the pleura. Diagnostic thoracentesis should be performed when a significant pleural effusion is present.

# Laboratory Evaluation/Antigen Detection

- ▶ Commercial assays can be used to detect capsular polysaccharide antigens of *S. pneumoniae* or *L. pneumophila* serogroup 1 in urine. The sensitivity of these tests is little affected by prior antibiotic treatment; indeed, results may remain positive several weeks after successful treatment. The degree of positivity for the *S. pneumoniae* urinary antigen test correlates with the *Pneumonia Severity Index* (PSI). The *S. pneumoniae* antigen test may also be applied on pleural fluid with a sensitivity and specificity of almost 100%.
- ▶ For *L. pneumophila* serogroup 1, the sensitivity is 60% to 80%, and the specificity is greater than 95%. Urinary antigen testing is currently the most helpful rapid test for the diagnosis of *Legionella* infections. The major limitation of urinary antigen tests is that currently available tests are intended to detect *L. pneumophila* serogroup 1 antigen only, although this is the most common cause of *Legionella* infection.
- ▶ Antigens for the many common respiratory viruses, influenza virus, respiratory syncytial virus, adenovirus, and parainfluenza viruses can be detected by direct immunofluorescence or by enzyme-linked immunoassay.



▶ **Serologic Evaluation**

- ▶ Before the development of nucleic acid amplification tests, serologic techniques were used to establish a microbiologic diagnosis for pneumonia caused by pathogens that cannot be readily cultured. Examples include common pathogens such as *M. pneumoniae*, *C. pneumoniae*, and *L. pneumophila*, and less common causes of pneumonia such as those caused by the agents of tularemia, brucellosis, and psittacosis, and certain viruses. Diagnosis usually requires that a convalescent specimen demonstrate a fourfold increase in *immunoglobulin (Ig) G* titer above that present in an acute specimen. These tests are not helpful in initial patient management but are of utility in defining the epidemiology of the pertinent infectious agents. Because IgM antibodies appear earlier than IgG antibodies, the detection of pathogen-specific IgM in serum has been used for the early serologic diagnosis of certain acute infections.

# Laboratory Evaluation/Nucleic Acid Amplification Tests


- ▶ Culture procedures for viruses and fastidious bacteria, *M. pneumoniae*, *C. pneumoniae*, *L. pneumophila*, and *Bordetella pertussis*, which normally do not colonize in the human respiratory tract, are too insensitive and too slow to be helpful in guiding therapy. These pathogens should be detected by nucleic acid amplification tests; their sensitivity is generally superior to that of the traditional procedures and some are considered as the “gold standard.”
- ▶ Real-time multiplex polymerase chain reaction assays detect respiratory viruses in both immunocompetent and immunosuppressed hosts.

# Invasive Diagnostic Techniques



- ▶ Bronchoscopic Samples
- ▶ Transthoracic Lung Aspiration

# Radiographic Evaluation

- ▶ Radiographic evaluation is necessary to establish the presence of pneumonia, because there is no combination of historical data, physical findings, or laboratory results that reliably confirms the diagnosis.
- ▶ The sensitivity of the chest radiograph is decreased in (1) patients with emphysema, bullae, or structural abnormalities of the lung, who may present with delayed or subtle radiographic changes; (2) obese patients, in whom it may be difficult to discern the existence of pneumonia; and (3) patients with very early infection, severe dehydration, or profound granulocytopenia.
- ▶ *Computed tomography* (CT) of the chest provides a more sensitive means of detecting minor radiographic abnormalities.







# Differential Diagnosis

Pulmonary edema

Pulmonary infarction

Acute respiratory distress syndrome

Pulmonary hemorrhage

Lung cancer or metastatic cancer

Atelectasis

Radiation pneumonitis

Drug reactions involving the lung


Extrinsic allergic alveolitis


Pulmonary vasculitis

Pulmonary eosinophilia

Organizing pneumonia

# Complications





# Treatment

- ▶ Risk stratify patients.
- ▶ Antibiotics.
- ▶ Treat complications

CURB-65	Clinical Feature	Points	
C	Confusion	1	
U	Urea > 7 mmol/L	1	
R	RR ≥ 30	1	
B	SBP ≤ 90 mm Hg OR DBP ≤ 60 mm Hg	1	
65	Age > 65	1	
CURB-65 Score	Risk group	30-day mortality	Management
0-1	1	1.5%	Low risk, consider home treatment
2	2	9.2%	Probably admission vs close outpatient management
3-5	3	22%	Admission, manage as severe

**TABLE 12. Antibiotic Therapy for Community-Acquired Pneumonia in Outpatients**

Risk Factors	Treatment
Previously healthy and no risk factor(s) for drug-resistant <i>Streptococcus pneumoniae</i>	Macrolide (azithromycin, clarithromycin, or erythromycin) or doxycycline
Risk factor(s) for drug-resistant <i>S. pneumoniae</i> or underlying comorbidities <sup>a</sup>	Respiratory fluoroquinolone (moxifloxacin, gemifloxacin, or levofloxacin) or $\beta$ -lactam <sup>b</sup> plus a macrolide or doxycycline


**TABLE 13. Empiric Antibiotic Therapy for Community-Acquired Pneumonia in Inpatients**

Inpatient Setting	Treatment
Medical ward	$\beta$ -lactam <sup>a</sup> plus a macrolide or doxycycline; or respiratory fluoroquinolone (moxifloxacin, gemifloxacin, or levofloxacin)
ICU	$\beta$ -lactam <sup>b</sup> plus either azithromycin or a fluoroquinolone <sup>c</sup> ; if penicillin allergic, a respiratory fluoroquinolone <sup>d</sup> plus aztreonam <sup>e</sup>
If risk factor(s) for <i>Pseudomonas aeruginosa</i> or gram-negative rods on sputum Gram stain	Antipseudomonal $\beta$ -lactam with pneumococcal coverage (cefepime, imipenem, meropenem, or piperacillin-tazobactam) plus ciprofloxacin or levofloxacin (750 mg); or antipseudomonal $\beta$ -lactam with pneumococcal coverage plus an aminoglycoside plus azithromycin; or antipseudomonal <sup>e</sup> $\beta$ -lactam with pneumococcal coverage plus an aminoglycoside plus a respiratory fluoroquinolone
If risk factor(s) for CA-MRSA, cavitary infiltrates, or compatible sputum Gram stain	Add vancomycin or linezolid to $\beta$ -lactam <sup>b</sup> plus either azithromycin or a fluoroquinolone <sup>c</sup>


# TYPICAL COMMUNITY ACQUIRED PNEUMONIA

ACUTE INFECTION OF THE LUNG PARENCHYMA  
ACQUIRED OUTSIDE OF THE HOSPITAL OR LESS  
THAN 48 HOURS AFTER HOSPITAL ADMISSION

FEVER, COUGH, SPUTUM PRODUCTION, RIGORS,  
PLEURITIC CHEST PAIN, DYSPNEA, TACHYCARDIA




MOST COMMON  
CAUSE OF TYPICAL CAP IS  
STREPTOCOCCUS PNEUMONIAE



BRONCHIAL BREATH  
SOUNDS AND CRACKLES  
ON AUSCULTATION

SPECIAL FINDINGS DUE TO  
LOBAR CONSOLIDATION INCLUDE:

- EGOPHONY (E TO A)
- WHISPERED PECTORILOQUY
- TACTILE FREMITUS (INCREASED)



INFILTRATE ON  
CHEST RADIOGRAPH

# Causes of Nonresponding Pneumonia

## INFECTIOUS

1. Resistant microorganisms
2. Community-acquired pneumonia (e.g., *Streptococcus pneumoniae*, *Staphylococcus aureus*)
3. Nosocomial pneumonia (e.g., *Acinetobacter*, methicillin-resistant *Staphylococcus aureus*, *Pseudomonas aeruginosa*)
4. Uncommon microorganisms (e.g., *Mycobacterium tuberculosis*, *Nocardia* spp., fungi, *Pneumocystis jirovecii*)
5. Complications of pneumonia
6. Empyema
7. Abscess or necrotizing pneumonia
8. Metastatic infection

## NONINFECTIOUS

1. Neoplasms
2. Pulmonary hemorrhage
3. Pulmonary embolism
4. Sarcoidosis
5. Eosinophilic pneumonia
6. Pulmonary edema
7. Acute respiratory distress syndrome
8. Organizing pneumonia
9. Drug-induced pulmonary disease
10. Pulmonary vasculitis

# Prevention of Pneumonia

- ▶ **Vaccines**

Prevention of pneumonia may be achieved by administering the influenza and pneumococcal vaccines.

- ▶ Smoking cessation.

## Recommendations for Administration of Influenza Vaccine\*

- ▶ Inactivated vaccine: All persons aged 6 months and older including pregnant women
- ▶ Live attenuated vaccine: Healthy, nonpregnant women aged 2 to 49 years without high-risk medical conditions

Risk Group	PCV13 Recommended	PPSV23 Recommended	PPSV23 Revaccination 5 Years after First Dose
Immunocompetent	<ol style="list-style-type: none"> <li>1.Cerebrospinal fluid leak</li> <li>2.Cochlear implant</li> </ol>	<ol style="list-style-type: none"> <li>1.Chronic heart diseases<sup>†</sup></li> <li>2.Chronic lung diseases<sup>‡</sup></li> <li>3.Diabetes mellitus</li> <li>4.Chronic liver diseases</li> <li>5.Cerebrospinal fluid leak</li> <li>6.Cochlear implant</li> <li>7.Alcohol</li> <li>8.Smoking</li> </ol>	
Asplenia	<ol style="list-style-type: none"> <li>1.Sickle cell disease/ hemoglobinopathy</li> <li>2.Congenital or acquired asplenia</li> </ol>	<ol style="list-style-type: none"> <li>1.Sickle cell disease/ hemoglobinopathy</li> <li>2.Congenital or acquired asplenia</li> </ol>	<ol style="list-style-type: none"> <li>1.Sickle cell disease/ hemoglobinopathy</li> <li>2.Congenital or acquired asplenia</li> </ol>
Immunocompromised	<ol style="list-style-type: none"> <li>1.HIV</li> <li>2.Congenital or acquired immunodeficiency</li> <li>3.Chronic renal failure</li> <li>4.Leukemia/lymphoma</li> <li>5.Generalized malignancy</li> <li>6.Solid organ transplant</li> <li>7.Multiple myeloma</li> <li>8.Iatrogenic immunosuppression<sup>§</sup></li> </ol>	<ol style="list-style-type: none"> <li>1.HIV</li> <li>2.Congenital or acquired immunodeficiency</li> <li>3.Chronic renal failure</li> <li>4.Leukemia/lymphoma</li> <li>5.Generalized malignancy</li> <li>6.Solid organ transplant</li> <li>7.Multiple myeloma</li> <li>8.Iatrogenic immunosuppression</li> </ol>	<ol style="list-style-type: none"> <li>1.HIV</li> <li>2.Congenital or acquired immunodeficiency</li> <li>3.Chronic renal failure</li> <li>4.Leukemia/lymphoma</li> <li>5.Generalized malignancy</li> <li>6.Solid organ transplant</li> <li>7.Multiple myeloma</li> <li>8.Iatrogenic immunosuppression</li> </ol>

QUESTIONS?