

BREAST MILK AND INFANT NUTRITION




Fareed Khdair, MD

Associate Professor

Chief, Section of Pediatric Gastroenterology,

Hepatology, and Nutrition

University of Jordan – School of Medicine





Topics

- Breast milk
 - Infant formulas
 - Failure To Thrive FTT
- 

Teaching modules in pediatric GI

- Web link:
- <http://radtf.indyrad.iupui.edu/radtf>



Home

About Us ▾

Departments ▾

Programs ▾

Facilities ▾

Achievements

Miscellaneous ▾

Contact Us

الموقع بالعربية



Anatomy and
Histology

Physiology and
Biochemistry

Pathology and
Microbiology and
Forensic Medicine

Pharmacology

Family and
Community Medicine

Internal Medicine

Radiology



Faculty of Medicine

"The faculty of medicine is the first medical school in Jordan. We are looking forward to being the first and most distinguished in the region."

The University of Jordan :: An

medicine.ju.edu.jo/Departments/School_DepthHome.aspx?deptname=Pediatrics

how to show hidden infected usb files

Employee Portal | Students Portal | Library | عربي

THE UNIVERSITY OF JORDAN Qs

Search

School

Dept. Home

Strategy

People

Courses

Programs

Featured Research

Study Plans

Forms

f

Department of Pediatrics

Departments



Faculty of Medicine

Department of Pediatrics

Waiting for www.facebook.com...



EN 11:13 PM 7/1/2017

http://medicine.ju.edu.jo/De x

medicine.ju.edu.jo/Departments/school_DeptCourses.aspx?DeptName=Pediatrics

how to show hidden infected usb files

☆

📅

📌

🏠

⚙️

☰



THE UNIVERSITY OF
JORDAN Qs

Employee Portal | Students Portal | Library | عربي

Search x

School

Dept. Home

Strategy

People

Courses

Programs

Featured Research

Study Plans

Forms

f

🐦

Department of Pediatrics

Departments ▾

 Department's Courses

📄 Course Level : Bachelor

Course Title	Program	Course Outline
Pediatrics – 1	Doctor of Medicine (M.D)	5th 2016-2017 JU syllabus ped1.pdf
Pediatrics – 2	Doctor of Medicine (M.D)	6th 2016-2017 JU syllabus (pead 2).pdf



Faculty of Medicine
Department of Pediatrics

Read www.google.com



EN 11:11 PM 7/1/2017



مركز الاعتماد
والتأمين الجوده



The University of Jordan

Accreditation & Quality Assurance Center

Course Syllabus

Course Name: Pediatrics – 1

facilities, Skills Lab/ Library/ related hospital facilities.

Required equipment:

Skills lab manikins and equipment

Equipped lecture room with IT support

References:

Required book (s), assigned reading and audio-visuals:

Nelson Textbook of Pediatrics, 19th edition, by R. Kliegman et al.

Recommended books, materials, and media:

1. **Nelson Essentials of Pediatrics, 6th Edition, by K Markdante.**
2. **Zitelli Atlas of Pediatric Physical Diagnosis, 4th Edition.**
3. **Harriet Lane Handbook of Pediatrics, 17th Edition.**
4. **Smith's Recognizable Patterns of Human Malformations, 6th Edition.**
5. **Online modules in pediatric GI (created by Indiana University).**

<http://radtf.indyrad.iupui.edu/radtf> Username: mfeist Password: student



Department of Educational Technology

(Powered by **EDACTIC™** Created by [Mark S. Frank, M.D.](#))

This is the School of Medicine's EDACTIC Server. Please log on.

Your UserID: Your Password:

(To re-initialize your session, [Click here](#))

...or click to a **public-access** site within our server:

1. Testing & Demo

This site is powered by **EDACTIC™** (Web Delivery Engine, Version 3.0.48 built 08/29/05 12:29:11)
EDACTIC stands for Education Delivered and Composed Through Internet Communication.
EDACTIC is a technology composed of software tools for authoring and delivering educational content on the Internet.

Note: This EDACTIC Enterprise Web Server application is licensed to IU School of Medicine to host web sites pertaining to *Medical Education*. Hosting content for other purposes or for any individual, group, or organization not affiliated with IU School of Medicine is prohibited.



Mark Feist MD

Login name : mfeist
Password : student



Indiana University School of Medicine Department of Educational Technology


(Powered by **EDACTIC™** Created by [Mark S. Frank, M.D.](#))

Welcome to Dr. Mark Feist's Educational Website

This website contains case-based modules which are part of a curriculum used to teach residents about pediatric gastroenterology, hepatology, and nutrition. The content of this website and the curriculum is based on the needs of primary care providers faced with children who present with gastrointestinal complaints.

The multiple choice and true/false questions in the modules are the means of teaching you much of the information. Do not get discouraged if you don't know the answers; you are not expected to know all the answers as this is the first time many of you have been exposed to this information. The incorrect answers on the multiple choice questions usually have an explanation of why they are incorrect and give you a little more information about that topic; therefore, clicking on all of answers will maximize your educational experience.

Click on the link below to access the modules.



[GI CASES](#)

Are you finished? Don't forget to [log off!](#)

All content on this Website is copyright © by IU School of Medicine.
Images may be reused for *educational purposes only*. All copyright watermarks must be left intact.

Powered by **EDACTIC™** -- Empowering Education Everywhere.



Indiana University School of Medicine Department of Educational Technology

(Powered by **EDACTIC™** Created by [Mark S. Frank, M.D.](#))

8 items were found.

Shown as...	Topic	Tests	Created	Updated
CaseStudy	Nutrition View		11/22/2005	02/23/2006
CaseStudy	Abdominal Pain View		09/27/2005	02/22/2006
CaseStudy	GI Bleeding View		07/25/2005	02/17/2006
CaseStudy	Diarrhea View		06/28/2005	06/14/2006
CaseStudy	Liver Disease II View		12/22/2004	02/24/2006
CaseStudy	Liver Disease I View		08/16/2004	01/24/2006
CaseStudy	Gastroesophageal Reflux View		07/14/2004	06/05/2006
CaseStudy	Constipation View		05/05/2004	08/23/2006

[Do Another Search](#) || [Home](#)





Indiana University School of Medicine Department of Educational Technology

(Powered by **EDACTIC™** Created by [Mark S. Frank, M.D.](#))

Pg 1 of 14 (GoTo: [Home](#) || [1](#) || [Next\(2\)](#) || [2](#) || [3](#) || [4](#) || [5](#) || [6](#) || [7](#) || [8](#) || [9](#) || [10](#) || [11](#) || [12](#) || [13](#) || [14](#))

Nutrition

Learning Objectives:

At the end of this module you should be able to:

- ▶ Describe the composition of various infant formulas and list indications for their usage
- ▶ Choose an infant formula for different clinical situations based on the protein, carbohydrate, and fat content of the formula
- ▶ Describe normal infant nutritional requirements
- ▶ List the most common causes of failure to thrive
- ▶ Describe the different types of failure to thrive based on trends in weight, height, and head circumference
- ▶ Assess nutritional status and diagnose malnutrition
- ▶ Describe the effects of pancreatic insufficiency on nutritional status
- ▶ Describe the risks that can be associated with nutritional rehabilitation
- ▶ Identify risk factors for and the presentation of deficiencies or toxicities of various vitamins and minerals

This is page 1 of 14

GoTo: [Top](#) || [1](#) || [Next\(2\)](#) || [2](#) || [3](#) || [4](#) || [5](#) || [6](#) || [7](#) || [8](#) || [9](#) || [10](#) || [11](#) || [12](#) || [13](#) || [14](#)

next page

[Home Page](#) || [Search](#) || [Search Hits](#)



Secure Search



McAfee

EN ?



6:55 PM
11/25/2012



Indiana University School of Medicine Department of Educational Technology

(Powered by **EDACTIC™** Created by Mark S. Frank, M.D.)

Pg 2 of 14 (GoTo: [Home](#) || [Prev\(1\)](#) || [2](#) || [Next\(3\)](#) || [4](#) || [5](#) || [6](#) || [7](#) || [8](#) || [9](#) || [10](#) || [11](#) || [12](#) || [13](#) || [14](#))

History: A 2-month-old male infant presents to your office because of irritability. The mother reports that the patient has become more irritable over the past few weeks and has begun to spit up after most feeds. The irritability occurs around the clock, and the patient is not sleeping well. The emesis is non-bilious, non-bloody and described as undigested formula. It usually occurs within a few minutes of completing the feeding. Mom estimates the amount of emesis as half of the volume consumed. She also states that he has a red, scaly rash on his arms, legs, and face which has gotten worse over the past few weeks.

Click on the links below for more history.

[Past Medical History](#)

[Family History](#)

[Social History](#)

[Diet History](#)

prev. page	This is page 2 of 14 GoTo: Top Prev(1) 2 Next(3) 4 5 6 7 8 9 10 11 12 13 14	next page
----------------------------	--	---------------------------

[Home Page](#) || [Search](#)

All content on this Website is copyright © by IU School of Medicine.
Images may be reused for *educational purposes only*. All copyright watermarks must be left intact.



Indiana University School of Medicine Department of Educational Technology

(Powered by **EDACTIC™** Created by [Mark S. Frank, M.D.](#))

Pg 4 of 14 (GoTo: [Home](#) || [1](#) || [2](#) || [Prev\(3\)](#) || [4](#) || [Next\(5\)](#) || [6](#) || [7](#) || [8](#) || [9](#) || [10](#) || [11](#) || [12](#) || [13](#) || [14](#))

🔍 **The most likely cause of this patient's symptoms is**

- ☐ A physiologic infantile gastroesophageal reflux
- ☐ B congenital alactasia (lactose intolerance)
- ☐ C cow's milk protein allergy
- ☐ D malrotation with midgut volvulus

🔍 **The treatment of choice for this patient would be a trial of**

- ☐ A Nutramigen
- ☐ B Isomil
- ☐ C Neocate
- ☐ D Enfamil Gentlease
- ☐ E goat's milk

🔍 **Response to therapy in this condition is usually seen**

- ☐ A immediately
- ☐ B within 12 hours
- ☐ C after 48-72 hours
- ☐ D after 2 weeks



This is page 4 of 14

GoTo: [Top](#) || [1](#) || [2](#) || [Prev\(3\)](#) || [4](#) || [Next\(5\)](#) || [6](#) || [7](#) || [8](#) || [9](#) || [10](#) || [11](#) || [12](#) || [13](#) || [14](#)



[Home Page](#) || [Search](#)

All content on this Website is copyright © by IU School of Medicine.
Images may be reused for **educational purposes only**. All copyright watermarks must be left intact.



Secure Search



McAfee

EN ?



7:53 PM

11/25/2012



Indiana University School of Medicine Department of Educational Technology

(Powered by **EDACTIC™** Created by [Mark S. Frank, M.D.](#))

8 items were found.


Shown as...	Topic	Tests	Created	Updated
CaseStudy	Nutrition View		11/22/2005	02/23/2006
CaseStudy	Abdominal Pain View		09/27/2005	02/22/2006
CaseStudy	GI Bleeding View		07/25/2005	02/17/2006
CaseStudy	Diarrhea View		06/28/2005	06/14/2006
CaseStudy	Liver Disease II View		12/22/2004	02/24/2006
CaseStudy	Liver Disease I View		08/16/2004	01/24/2006
CaseStudy	Gastroesophageal Reflux View		07/14/2004	06/05/2006
CaseStudy	Constipation View		05/05/2004	08/23/2006

[Do Another Search](#) || [Home](#)



BREAST MILK

Components of breast milk





Benefits of breast milk

- Immunity and protection against infection:
 - Antibodies in the milk
 - Cytokines
 - Normal flora growth factors
- 



Benefits of breast milk

- Available 24/7
 - Cheaper!
 - No need for preparation : bottle/ water
- 



Benefits of breast milk

- Better intelligence
 - Protects against obesity
- 


Table 42-3 CONDITIONS FOR WHICH HUMAN MILK HAS BEEN SUGGESTED TO HAVE A PROTECTIVE EFFECT

Acute disorders

- Diarrhea
- Otitis media
- Urinary tract infection
- Necrotizing enterocolitis
- Septicemia
- Infant botulism

Chronic disorders

- Insulin-dependent diabetes mellitus
- Celiac disease
- Crohn's disease
- Childhood cancer
 - Lymphoma
 - Leukemia
- Recurrent otitis media
- Allergy
- Obesity and overweight
- Hospitalizations
- Infant mortality



Adapted from the Schanler RJ, Dooley S: *Breastfeeding handbook for physicians*, Elk Grove Village, IL, 2006, American Academy of Pediatrics.

Mother- Baby Bonding



Benefits of breast milk to the mother

- Helps with mother weight loss
- Helps uterine contraction post delivery
- Helps with to get rid of pregnancy hormones
- Act as contraceptive : Lactation induced amenorrhea LAM



BREAST IS BEST

Patterns of milk supply

Table 42-5 PATTERNS OF MILK SUPPLY

DAY OF LIFE	MILK SUPPLY
Day 1	Some milk (~5 mL) may be expressed
Days 2-4	Lactogenesis, milk production increases
Day 5	Milk present, fullness, leaking felt
Day 6 onward	Breasts should feel “empty” after feeding

Adapted from Neifert MR: Clinical aspects of lactation: promoting breastfeeding success, *Clin Perinatol* 26:281–306, 1999.

How to Encourage breast feeding

Table 42-1 STEPS TO ENCOURAGE BREAST-FEEDING IN THE HOSPITAL: UNICEF/WHO BABY-FRIENDLY

HOSPITAL INITIATIVES

Provide all pregnant women with information and counseling
Document the desire to breast-feed in the medical record
Document the method of feeding in the infant's record
Place the newborn and mother skin-to-skin, and initiate breast-feeding within 1 hr of birth
Continue skin-to-skin contact at other times and encourage rooming in
Assess breast-feeding and continue encouragement and teaching on each shift

MOTHERS TO LEARN

Proper position and latch on
Nutritive sucking and swallowing
Milk production and release
Frequency and feeding cues
Expression of milk if needed
Assessment of the infant's nutritional status
When to contact the clinician

ADDITIONAL INSTRUCTIONS

Refer to lactation consultation if any concerns arise
Infants should go to the breast at least 8-12 times/24 hr day and night
Avoid time limits on the breasts; offer both breasts at each feeding
Do not give sterile water, glucose, or formula unless indicated
If supplements are given, use cup feeding, a Haberman feeder, fingers, or syringe feedings
Avoid pacifiers in the newborn nursery except during painful procedures
Avoid antilactation drugs

UNICEF, United Nations Children's Fund; WHO, World Health Organization.

Table 42-4 ABSOLUTE AND RELATIVE CONTRAINDICATIONS TO BREAST-FEEDING DUE TO MATERNAL HEALTH CONDITIONS

MATERNAL HEALTH CONDITIONS	DEGREE OF RISK
HIV and HTLV infection	In the USA, breast-feeding is contraindicated In other settings, health risks of not breast-feeding must be weighed against the risk of transmitting virus to the infant
Tuberculosis infection	Breast-feeding is contraindicated until completion of approximately 2 wk of appropriate maternal therapy
Varicella-zoster infection	Infant should not have direct contact to active lesions Infant should receive immune globulin
Herpes simplex infection	Breast-feeding is contraindicated with active herpetic lesions of the breast
CMV infection	May be found in milk of mothers who are CMV seropositive Transmission through human milk Causing symptomatic illness in term infants is uncommon.
Hepatitis B infection	Infants routinely receive hepatitis B immune globulin and hepatitis B vaccine if mother is HbsAg positive No delay in initiation of breast-feeding is required
Hepatitis C infection	Breast-feeding is not contraindicated
Alcohol intake	Limit maternal alcohol intake to <0.5 g/kg/day (for a woman of average weight, this is the equivalent of 2 cans of beer, 2 glasses of wine, or 2 oz of liquor)
Cigarette smoking	Discourage cigarette smoking, but smoking is not a contraindication to breast-feeding
Chemotherapy, radiopharmaceuticals	Breast-feeding is generally contraindicated

CMV, cytomegalovirus; HbsAg, hepatitis B surface antigen; HIV, human immunodeficiency virus; HTLV, human T-lymphotropic virus.



INFANT FORMULAS

Infant formula

- Can be classified according to their content:
- Protein content
- Carbohydrate content
- Fat content



How to think about formulas

Formula type	Prtn content	CHO content	fat content

Infant Formulas – Protein Content

- Divided into 4 classes of formulas
 - Cow's milk based formulas
 - variable prtn content
 - variable whey: casein ratio
 - Soy formulas
 - Casein **hydrolysate** formulas
 - Extensive **Vs** partial
 - Amino acid based formulas



When do we have to change the formula class?

- When you suspect

Cow's milk protein allergy



Cow's milk protein allergy

- It is a clinical diagnosis
- Any combination of the following
- Vomiting
- Abd distension
- Diarrhea
- Blood in stool
- Irritability and fussiness
- eczema



in a formula fed or breast fed infant



Indiana University School of Medicine
Department of Educational Technology

(Powered by **EDACTIC™** Created by Mark S. Frank, M.D.)

Back Button

Protein Content of Infant Formulas			
Protein Source	Examples	Indications	Price*
Cow's Milk	Enfamil with Iron	Normal GI tract; Enfamil AR used for gastroesophageal reflux	\$
	Enfamil Lipil		
	Similac with Iron		
	Similac Advance		
	Carnation Good Start		
	Good Start Supreme		
	Enfamil Gentlease		
	Enfamil AR		
	Store Brands		
Soy	Prosobee	Cow's milk protein allergy, Lactose malabsorption, or Galactosemia	\$
	Isomil		
	Alsoy		
	Store Brands		
Casein Hydrolysate	Nutramigen	Cow's milk and/or soy allergy; Alimentum and Pregestimil are also used for malabsorption	\$\$
	Alimentum Pregestimil		
Amino Acids	Neocate Elecare	Severe protein allergy not responsive to casein hydrolysate formula	\$\$\$-\$\$\$\$

Bebelac HA
Nan HA

S26
Nan
Bebelac
Seha
Similac
Ronlac
AR formulas
"Sensitive" / LF/

* Each \$ = approximate cost of standard cow's milk based formula

As you move down this table from cow's milk to soy to hydrolysate to amino acid based formulas, the formulas become less antigenic; formulas within a class are similarly antigenic to one another. When choosing a formula to treat milk protein allergy, you should progress down the table. It is not beneficial to change to a different formula



Secure Search



McAfee

Infant Formulas – Carbohydrate Content


- Main types of carbohydrates in formulas
 - Lactose
 - Sucrose
 - Glucose polymers :
- What type of formula should be used in patients with galactosemia? Why?
 - formulas that do not contain lactose
- Which formulas contain sucrose?
 - Alimentum and soy formulas except Prosobee
- Glucose polymers :
 - Hydrolysate and AA based formulas



Infant Formulas – Fat Content

- Main types of fats in formulas
 - Long chain triglycerides (LCTs)
 - Medium chain triglycerides (MCTs)
- 

Absorption of MCT vs LCT



- When are MCTs beneficial?
 - Impaired fat absorption or lymphatic abnormalities
- Which formulas contain MCTs?
 - Alimentum (33%), **Pregestimil (55%)**, Alfare 38%
 - Elecare (33%)
 - Portagen (87%), Vital HN (45%)

Proper preparation of formula

- Hand washing
- **READ THE INSTRUCTIONS**
- Most formulas in the local market:
 - 1 scoop powder in 30 ml
(nan, babelac, seha, ronlac)
 - Except : S26, similac 1 scoop in 60 ml

Calorie content of formulas

- Regular formula (breast milk)
- 67 kcal : 100 ml
- 20 Kcal : 30 ml (oz) 20 Kcal/ oz

Regular calorie needs for infant

- At least 100 kcal/ kg/ day
- We need to always calculate the calorie intake from formula
- $\text{Total volume} / \text{weight} \times \text{formula calorie concentration} = \text{Kcal} / \text{kg} / \text{day}$

Example

- 1 month old infant
 - 4 kg
 - Regular Formula intake 90 ml q 3 hrs
 - What is the daily total caloric intake ?
-
- Total volume / weight X formula Conc.
 - $(90 \times 8) / 4 \times 67/100$
= 120 kcal / kg/ day



Failure To Thrive

FTT






Failure To Thrive

FTT

- The inability to maintain the **expected rate** of growth over time.
 - Growth is assessed by plotting the patient's growth parameters over **subsequent visits** and comparing the growth rate to normal population growth rates for age.
- 


- 
- One set of measurements can not assess rate of growth and therefore is not sufficient to diagnose failure to thrive
- 

Failure to Thrive




- List the three main causes of this type of growth pattern
- Type I failure to thrive
 - Inadequate caloric intake
 - Excessive loss of calories
 - Increased metabolic demands

Failure to Thrive



- List three causes of this type of growth pattern
- Type II failure to thrive
 - Constitutional growth delay
 - Genetic short stature
 - Hypothyroidism
 - Growth hormone deficiency
 - Hypopituitarism
 - Chronic malnutrition

Failure to Thrive



- List three causes of this type of growth pattern
- Type III failure to thrive
 - Congenital infections
 - Chromosomal abnormalities
 - Prenatal exposure to toxins

Type I Failure to Thrive

- Inadequate caloric intake
 - Inappropriate feeding regimen/schedule
 - Formula prepared incorrectly
 - Decreased appetite or feeding dysfunction/refusal
- Excessive loss of calories
 - GER or vomiting
 - Diarrhea/malabsorption
- Increased metabolic demands
 - Hyperthyroidism, diencephalic syndrome



Nutrition assessment

- At EMR
- 

Jordan University Hospital
Window Help Tools Security View Object File

Admission Note Ped

Patient Profile
nutritional scr & functional assessment
PHYSICAL EXAMINATION
plan of care

p_id 1172503 adm 2 حلا ياسر المصري Admission Date 08-10-2019

age 0 Gender أنثى Admitted Via ED OPD طباعة

History was taken from : Patient Others استفسار

Chief Complaint failure to gain weight(FTT)

Present Illness 6 month old female syrian baby,Down syndrome ,lives in مخيم الزعتري admitted as a case of FTT (type 3).

Antenatal Hx mentioned

Natal Hx & Birth Hx mentioned

Postnatal Hx mentioned

Nutritional Hx mentioned

Developmental mentioned

Vaccination mentioned

Past Medical Hx: 3 previous admission

Past Surgical Hx mentioned

Allergy: NKDA

Social Hx: no pets at home ,no smoking at home

Family Hx cons (1st degree)

Family tree mentioned

Diet pediasure,90m Q3hrs

REVIEW OF SYSTEM(Give details and negative)

Head & Neck mentioned

ENT: mentioned

Cardiac/respiratory mentioned

GI: mentioned

Genitourinary: mentioned

Neurological: mentioned

Musculoskeletal: mentioned

Psychiatry

Others

Patient Profile

nutritional scr & functional assessment

PHYSICAL EXAMINATION

plan of care

weight Height BMI Normal 14_25Oral diet prescribed: enteral feeding oral supplements infantrene

one (1) point

☐ Diabetes☒ GI mobility or absorption disorder☐ critical care patient☐ Chronic renal failure☐ severe anemia-abnormal labs HGB<8.0☐ special diet(specify) ☐ Diarrhea lasting more than 3 Weeks☐ Nausea or vomiting more than 3days per week for more than 1 monthlabel

Weight Loss

☐ pressure ulcer /wounds☐ Chewing difficulties☐ impaired swallowing☐ Major burns☐ lab:albuminb less than 3.0☐ Pitting edema(3+-4+)**Functional assessment criteria**1.Need assistant in performing activity of daily living(walking,feeding,grooming ,driving,cleaing,toileting,shopping,_) ☐ yes ☒ no2.presence of neurological deficit (weakness,paralysis,paresthesia,imbalance) ☐ yes ☒ no3.presence of muscle skeletal problem(arthritis ,prosthesis fractures,joint replacement) ☐ yes ☒ no4.Limited range of motion (stiffness,post-surgery) ☐ yes ☒ no5.Presence of pain interfere with function ☐ yes ☒ no6.Amputation ☐ yes ☒ nolabel

Presence of any positive(yes)answer indicates a need for further functional assessment by specialist

Vital signs

Temp	Pulse	Resp.	BP	Hand circumference	Pain score	Weight	Height
37.3	150	31			0	3.3	53

Nutrition assessment

Visit date:	File no:	Name:		
Date of birth:	Gender:	Age:	Percentiles wt:	Percentiles ht:
Previous wt:	Current wt:	Ht:	Hip Circumference:	
BMI:	MUAC:	Waist Circumference:		
Reason for the Visit:				
Family History:				
Diet History:				
Dietary Recall:				
Like Food:	Dislike Food:	Allergies :	PA: (Sedentary / Low Active/ Active / Very Active)	
Fast Food:	No. of Meals:	Fruits and Veg. intake:		
Duration of Screen Viewing: ☐ Hour (s)				
Patient's Calories intake:		His /Her Energy Req.:		
Lab Results:				
Assessment:				
Plan and Recommendations :				
*Diabetic Patient:				
Carbs Count Method:	☐ Grams	☐ Serving		
ICR:	ISF:	Type of insulin: (MDI, Mixtard)		
No. of Gluco-check Readings:		Insulin dose determined by: (Mother, Patient, Other)		
Insulin dose determined: (Estimation, Precisely by grams, Serving size)				
Notes:				
*Ketogenic Diet				
Energy Req.=	Fat 80%=	Pro 15%=	CHO 5%=	
Notes:				



Nutritional Assessment

- History
 - Intake, losses, past growth, parental heights
 - Anthropometrics
 - Height/length, weight, head circumference, BMI
 - Skinfold thickness, mid-upper arm circumference
 - Physical Exam
 - Decreased fat stores, muscle wasting, edema
 - Lab
 - Visceral proteins, CBC, K, mag, phos, zinc
- 

Nutritional Status

- Wasting

- Weight:length ratio or BMI $<3^{\text{rd}}$ percentile
- Often seen in type I failure to thrive
- Indicative of acute malnutrition
- Typically responds to nutritional support

- Stunting

- Height $<3^{\text{rd}}$ percentile for age
- Often have a normal weight:length ratio or BMI
- Chronic malnutrition may progress to stunting

Nutritional Rehabilitation

- How do you decide between enteral and parenteral support?
 - Use parenteral route when, and only when, enteral support is not possible or not adequate to meet the nutritional needs of the patient
- What type of enteral support should you use?
 - Use most physiologic method tolerated by patient
 - Most physiologic to least physiologic:
 - Increasing caloric density → oral supplements → gastric bolus → gastric continuous → jejunal continuous

Complications of Nutritional Support

- What are risk factors for developing the refeeding syndrome?
 - Moderate to severe malnutrition
- What are the laboratory findings?
 - Hypokalemia, hypomagnesemia, and hypophosphatemia
- How do you avoid this complication?
 - Advance feedings and/or TPN slowly
 - Carefully monitor and supplement K, Mag, Phos



Complications of Nutritional Support

- Discuss complications that may be seen with enteral support
 - Tube malposition
 - Irritation or infection of tube site
 - Discuss complications that may be seen with parenteral support
 - Infection
 - Metabolic derangements
 - Mechanical complications
- 

THE END

QUESTIONS?

